Finite field- and BCH codes

2^{nd} December 2005

Definition 1. Let G be a group. Then a *coset* is a subgroup H of G which is either a *left coset* of H, that is $xH = \{xh : h \in H\}$ for some x in G, or a *right coset* $Hx = \{hx : h \in H\}$ of the same.

§

Definition 2. Let polynomials $f_1(x), \ldots, f_r$ in $\mathbf{F}_q[x]$ be non-zero. Then the *least common multiple* $\operatorname{lcm}(f_1(x), \ldots, f_r(x))$ of $f_1(x), \ldots, f_r(x)$ is the monic polynomial of the lowest degree which is a multiple of all $f_i(x)$, $i = 1, \ldots, r$.

§

Problem 1. Prove that for non-zero polynomials $f_1(x), \ldots, f_r(x)$ in $\mathbf{F}_q[x]$,

$$\operatorname{lcm}(f_1(x), \ldots, f_r(x)) = \operatorname{lcm}(\operatorname{lcm}(f_1(x), \ldots, f_{r-1}(x)), f_r(x))$$

§

Note 1. Let $f_1(x), \ldots, f_r(x)$ in $\mathbf{F}_q[x]$ have the factorisations,

$$\begin{aligned} f_1(x) &= a_1 (p_1(x))^{e_{11}} \cdots (p_n(x))^{e_{1n}} \\ &\vdots \\ f_r(x) &= a_r (p_1(x))^{e_{r1}} \cdots (p_n(x))^{e_{rn}} \end{aligned}$$

where $a_1, \ldots, a_r$ are in $\mathbf{F}_q^*$, $e_{ij} \geq 0$, and $p_i(x)$ are distinct monic irreducible polynomials over $\mathbf{F}_q$, then

$$\operatorname{lcm}(f_1(x), \ldots, f_r(x)) = (p_1(x))^{\max(e_{11}, \ldots, e_{r1})} \cdots (p_n(x))^{\max(e_{1n}, \ldots, e_{rn})}$$

§

Theorem 1. Let $f(x), f_1(x), \ldots, f_r(x)$ be polynomials over $\mathbf{F}_q$. If $f(x)$ is divisible by every polynomial f_i, for $i = 1, \ldots, r$, then $f(x)$ is also divisible by $\operatorname{lcm}(f_1(x), \ldots, f_r(x))$.

Proof. Consider first the case where there are only two different polynomials, $f_1(x)$ and $f_2(x)$. The prime components of $f_1(x)$ and $f_2(x)$ may be grouped into those which are unique among them and those which are shared. Since $f(x) = u_1(x)f_1(x) + r_1(x)$ and $f(x) = u_2(x)f_2(x) + r_2(x)$, it follows that $f(x)$ contains both of these two groups of primes. In other words, $f(x) = u(x)\operatorname{lcm}(f_1(x)f_2(x)) + r(x)$.
Next, consider the case where there are more than two f_i's. Suppose for $f(x)$, that $f(x) = u_r(x)\operatorname{lcm}(f_1(x), \ldots, f_r(x))$. Then if we let $f_c(x) = \operatorname{lcm}(f_1(x), \ldots, f_r(x))$, and if we introduce another polynomial $f_{r+1}(x)$ such that $f(x) = u_{r+1}f_{r+1} + r_{r+1}(x)$, then following the same line of reasoning as the above we have,

$$\operatorname{lcm}(f_1(x), \ldots, f_{r+1}(x)) | f(x)$$

¶

Definition 3. A non-empty subset S of a ring R is called a *subring* of R if the elements of S form a ring with respect to the operations defined in R.

§

Theorem 2. Let R be a ring. Then a non-empty subset S of R is a subring if and only if S is closed under addition, multiplication, and the formation of additive inverse.

Proof. Since S is a subset of R, additive associativity, identity and commutativity are inherited to S from R. The existence of the inverse for each element s in S is certain provided that the formation of an additive inverse is guaranteed. And similarly in the case of multiplication, both associativeness and distributiveness hold once we know that S is closed under multiplication. ¶

Definition 4. Let R be a ring. We call an *ideal* in R a sub ring I having such property that for all i in I, then both xi and ix are also in I for every element x in R. Further, if I is a proper subset of R, then it is called a *proper ideal*. By *trivial ideal* one means either the *zero ideal* $\{0\}$ consisting of the zero element alone, or the ring R itself.

§

Note 2. The significance of the ideals in a ring is that they let us construct other rings from the first. The cosets of a ring R is a partition of R into equivalence sets, which are non-empty and disjoint, the union of which is the whole of the ring R.

§

Definition 5. Let R be a ring and I an ideal in it. Then two elements x and y in R are said to be *congruent modulo* I, denoted by $x \equiv y (\mathrm{mod}\, I)$, if $x - y$ is in I. Since there is only ideal, we may a write this congruence as simply $x \equiv y$.

§

Note 3. The congruence modulo I of a ring R as defined in Definition 5 is an equivalence relation since it is true that $x \equiv x$ for every x; $x \equiv y$ implies $y \equiv x$; and $x \equiv y$ and $y \equiv z$ implies $x \equiv z$.

§

Note 4. Congruences can be added and multiplied as if they were ordinary equations. In other words, if $x_1 \equiv x_2$ and $y_1 \equiv y_2$, then $x_1 + y_1 \equiv x_2 + y_2$ and $x_1 y_1 \equiv x_2 y_2$.

§

Definition 6. Let R be a ring and let x be an element of R. Then the *coset* $[x]$ containing x is the set of all elements y such that $y \equiv x$. Then,

$$\begin{aligned}[x] = \{y : y \equiv x\} &= \{y : y - x \in I\} \\ &= \{y : y - x \in i \,\text{for some}\, i \in I\} \\ &= \{y : y = x + i \,\text{for some}\, i \in I\} \\ &= \{x + i : i \in I\} = x + I\end{aligned}$$

Furthermore, $[x] = [x_1]$ means that $x \equiv x_1$, that is to say, $x - x_1$ is in I. Here x and x_1 are called *representatives* of the coset which contains them.

§

Definition 7. A *quotient ring*, aka *residue-class-*, *factor-*, or *difference ring*, is a ring having the form of a quotient A/i of a ring A and one of its ideal i. In other words, the quotient ring of R with respect to I the ring $R/I = \{x + I : x \in R\}$, where $x + I = \{x + i : i \in I\}$ is the coset of an element x in R, and where addition and multiplication are defined as,

$$[x] + [y] = [x + y]$$

and

$$[x] \cdot [y] = [xy]$$

§

Theorem 3. The zero element of R/I is $0 + I = I$, the negative of $x + I$ is $(-x) + I$. If R is commutative, then R/I is also commutative. If R has an identity 1 and a proper ideal I, then R/I has an identity $1 + I$.

§

Problem 2. Prove Theorem 3.

§

Theorem 4. Let R be a ring and I an ideal of R. Then, for x and y in R,

$$(x + I) + (y + I) = (xy) + I$$

and

$$(x + I)(y + I) = xy + I$$

Proof. Let a and b be any two elements of the ideal I. Then,

$$(x+a)+(y+b) = x+a+y+b = (x+y)+(a+b) = (x+y)+p$$

where $p = a+b$ is in I. Further,

$$\begin{aligned}(x+a)(y+b) &= xy+bx+ay+ab \\ &= xy+c+d+e = xy+f\end{aligned}$$

where $c = bx$, $d = ay$, $e = ab$ and $f = c+d+e$ are all elements of I. ¶

Note 5. Theorem 4 and Note 4 show that the quotient ring R/I defined in Definitions 7 is independent of the choice of x and y in the cosets $x+I$ and $y+I$. In other words, the cosets $[x+y]$ and $[xy]$ resulted from addition and respectively multiplication in no ways depend on the particular representatives x and y chosen for the cosets $[x]$ and $[y]$ that go into them. This means that, if $x_1 \equiv x$ and $y_1 \equiv y$, then $[x_1+y_1] = [x+y]$ and $[x_1y_1] = [xy]$, or equivalently $x_1+y_1 \equiv x+y$ and $x_1y_1 \equiv xy$.

§

Example 1. Some examples of quotient ring are $\mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z}$ and $\mathbf{Z}_6 = \mathbf{Z}/6\mathbf{Z}$.

Theorem 5. The polynomial ring $F[x]$ is a commutative ring with identity.

Proof. $F[x]$ is a ring over the field F since under addition it is closed, associative and commutative, and has 0 as the identity and the inverse $-f(x)$, where $f(x) \in F[x]$; and under multiplication it is associative, distributive and commutative, and has 1 as the identity. ¶

Definition 8. Let R be a commutative ring with identity. Then for any a in R the *principal ideal* generated by a is $\langle a\rangle = aR = \{ar : r \in R\}$. Further, R is called *principal ideal ring* if all its ideals are of this form.

§

Theorem 6. Let F be a field. Then the polynomial ring $F[x]$ is a principal ideal ring.

Proof. The polynomial ring $F[x]$ being a commutative ring with identity, it remains only to show that all its ideals are of the form $\langle a\rangle\, R = aR = \{ar : r \in R\}$, where a is in R. Let I be an ideal of $F[x]$. If $I = 0$, then I is a principal ideal generated by 0. If $I \neq 0$, then choose $0 \neq f(x) \in I$ such that $\deg f \leq \deg g$ for all non-zero $g(x)$ in I. Write $g(x) = q(x)f(x) + r(x)$. If $\deg g < \deg f$, then $q = 0$ and $r = f$. On the other hand, if $n = \deg f \leq \deg g$, then either r is 0 or $\deg r < \deg f$. Let

$$f(x) = a_0x^n + \cdots + a_n$$

and

$$g(x) = b_0x^m + \cdots + b_m$$

Then, with $a_0 \neq 0$,

$$g(x) = a_0^{-1}b_0x^{m-n}f(x) + g_1(x) \tag{1}$$

where $\deg g_1 \leq m-1$. Then

$$g_1(x) = q_1(x)f(x) + r(x) \tag{2}$$

From this it follows that either $r = 0$ or $\deg r < \deg f$. From Equation's 1 and 2, $g(x) = q(x)f(x) + r(x)$, where $q(x) = a_0^{-1}b_0x^{m-n} + q_1$ is in $F[x]$. If $r \neq 0$, then $r(x)$ is in I and $\deg r < \deg f$, which contradicts our choice of $f(x)$. Therefore $g = qf$ and I is a principal ideal generated by $f(x)$. ¶

Definition 9. Let R be a commutative ring with identity. Then a non constant $f(x)$ in $R[x]$ is said to be *reducible* if, for some $g(x)$ and $h(x)$ in $R[x]$, $f(x) = g(x)h(x)$ implies either $\deg g(x) = 0$ or $\deg h(x) = 0$. Otherwise $f(x)$ is said to be *reducible*.

§

Theorem 7. Let F be a field $f(x)$ in $F[x]$ an irreducible polynomial. Then $F[x]/\langle f(x)\rangle$ is a field.

Proof. Let I be the ideal $\langle f(x)\rangle$ of $F[x]$ generated by $f(x)$. If $I = F[x]$, then $f(x)$ has an inverse, that is $1 = f(x)g(x)$ for some $g(x)$ in $F[x]$. Then $f(x)$ is a constant polynomial, which contradicts

our statement of the theorem. Therefore $F[x]/I$ has at least two elements, and $F[x]/I$ being a polynomial ring it is a commutative ring with identity. Let $g \in F[x]$ and $g \notin I$. Then,

$$J = \{a(x)f(x) + b(x)g(x) : a(x), b(x) \in F[x]\}$$

is an ideal of $F[x]$ and there exists $h(x)$ in $F[x]$ such that $J = \langle h(x) \rangle$. But $f(x) = 1f(x)+0g(x)$ is in J, and thus $f(x) = a(x)h(x)$ for some $a(x)$ in $F[x]$. The polynomial $f(x)$ being irreducible, either $\deg h(x) = 0$ or $\deg a(x) = 0$. If the latter is the case, then $a(x)$ is a unit in $F[x]$, and then $h(x)$ is in I, hence $J = I$, and hence a contradiction since we began with g being in J but not in I. Therefore it must be the case that $h(x)$ is a unit in $F[x]$, hence J is a unit, and thus $1 = a(x)f(x) + b(x)g(x)$ for some $a(x)$ and $b(x)$ in $F[x]$. And then $1 + I = I + b(x)g(x) = (b(x) + I)(g(x) + I)$. Thus $g(x) + I$ has an inverse and $F[x]/I$ is a field. ¶

Definition 10. Let K be a field and F a subfield of K. Then K is called an *extension* of the field F, denoted by $K|_F$. Since K has multiplication, it is a vector space over F. The dimension of the vector space K over F is called the *degree* $[K : F]$ of the extension K of F. The extension $K|_F$ is said to be *finite* if the degree $[K : F]$ is finite.

§

Definition 11. A *prime subfield* of a field F is the intersection of all subfields of F. It is the smallest of all subfields of F, and is unique. A *prime field* is a field which has no proper subfields.

§

Definition 12. Let $K|_F$ be an extension of a field F. Then $\alpha \in K$ is said to be *algebraic* over F if there exists $f(x)$ in $F[x]$ which has α as a root. Let α in K be algebraic over F and consider $A = \{f(x) \in F[x\} : f(\alpha) = 0]$. Here A is an ideal of the principal ideal domain $F[x]$. Let $m_1(x)$ in $F[x]$ be a generator of A. If a is the coefficient of the highest power of x in $m_1(x)$, then $m(x) = a^{-1}m_1(x)$ is a monic polynomial with $\deg m(x) = \deg m_1(x)$, and $m(x)$ is also a generator of A. Let $m(x) = r(x)s(x)$ for some $r(x)$ and $s(x)$ in $F[x]$. Then either $r(\alpha) = 0$ or $s(\alpha) = 0$, that is either $m(x)|r(x)$ or $m(x)|s(x)$. But $\deg m = \deg r + \deg s$, therefore either $\deg r(x) = 0$ or $\deg s(x) = 0$. Hence $m(x)$ is irreducible. Since $m(x)$ is monic, irreducible and is of the least degree possible while admitting α as a root, therefore $m(x)$ is called the *minimal polynomial* of α over $F[x]$.

§

Theorem 8. Let C be an (n, k) linear code over F_q with prith-check matrix H, and $d(C)$ the smallest number of column of H that are linearly dependent. Then if every subset of $2t$ or fewer columns of H is linearly independent, the code is capable of correcting all error patterns of weight $w \leq 1$.

Proof. When $q = 2$, linear independence amounts to summing to $\mathbf{0}$. The code words of C are those vectors $\mathbf{x}$ in $V_n(F_q)$ for which $H\mathbf{x}^T = \mathbf{0}$. But $H\mathbf{x}^T$ is a linear combination of the columns of H, that is to say, if $H = [\,\mathbf{c}_1 \quad \cdots \quad \mathbf{c}_n\,]$, then $H\mathbf{x}^T = x_1\mathbf{c}_1 + \cdots + x_n\mathbf{c}_n$. Hence a non-zero code word of weight w gives a nontrivial linear dependence among w columns of H, and vice versa. ¶

Corollary 8[1]. If $q = 2$ and all possible linear combinations of up to e columns are distinct, then $d(C) \geq 2e + 1$, and C can then correct all patterns of weight e or less.

Problem 3. Prove Corollary 8[1].

§

Note 6. Hamming codes correct single errors. An extension of this is to the Bose-Chaudhuri-Hocquenghem codes which could correct multiple errors. In the case of Hamming code of length $n = 2^m - 1$, the parity-check matrix is given by $H = [\,\mathbf{v}_0 \quad \ldots \quad \mathbf{v}_{n-1}\,]$, where $(\,\mathbf{v}_0 \quad \cdots \quad \mathbf{v}_{n-1}\,)$ is some ordering of the $2^m - 1$ non-zero column vectors in $V_m = V_m(F_2)$. The $m \times n$ matrix H takes m parity-check bits for the code to be able to correct one error. We may extend H such that it has m more rows and could correct two errors. Then,

$$H_2 = \begin{bmatrix} \mathbf{v}_0 & \cdots & \mathbf{v}_{n-1} \\ \mathbf{w}_0 & \cdots & \mathbf{w}_{n-1} \end{bmatrix}$$

where $\mathbf{w}_0, \ldots, \mathbf{w}_{n-1}$ are in V_m. Since $\mathbf{v}_i$'s are distinct, we may look at the mapping from $\mathbf{v}_i$ to $\mathbf{w}_i$ as a function from V_m into itself, then

$$H_2 = \begin{bmatrix} \mathbf{v}_0 & \cdots & \mathbf{v}_{n-1} \\ \mathbf{f}(\mathbf{v}_0) & \cdots & \mathbf{f}(\mathbf{v}_{n-1}) \end{bmatrix}$$

Then H_2 will define a code which corrects two errors if and only if the syndromes of the $1+n+\binom{n}{2}$ error patterns of weights 0, 1 and 2 are all distinct. Any such syndrome is a sum of a subset of columns of H_2, and therefore a vector in V_{2m}. Let the syndrome be $\mathbf{s} = (\, s_1 \quad \ldots \quad s_{2m}\,) = (\, \mathbf{s}_1 \quad \mathbf{s}_2\,)$, where $\mathbf{s}_1 = (\, s_1, \ldots, s_m\,)$ and $\mathbf{s}_2 = (\, s_{m+1}, \ldots, s_{2m}\,)$ are both in V_m. Defining $\mathbf{f}(\mathbf{0}, \mathbf{0}) = \mathbf{0}$ we consider a pair of errors occuring at i^{th}- and j^{th} position's, $\mathbf{s} = (\mathbf{v}_i + \mathbf{v}_j, \mathbf{f}(\mathbf{v}_i) + \mathbf{f}(\mathbf{v}_j))$. Then the system of equations,

$$\mathbf{u} + \mathbf{v} = \mathbf{s}_1$$
$$\mathbf{f}(\mathbf{u}) + \mathbf{f}(\mathbf{v}) = \mathbf{s}_2$$

has at most one solution $(\mathbf{u}, \mathbf{v})$ for each pair of vectors from V_m. By trial and error we may find neither the linear mapping $\mathbf{f}(\mathbf{v}) = T\mathbf{v}$ nor the nonlinear polynomial of degree 2 works, but $\mathbf{f}(\mathbf{v}) = \mathbf{v}^3$ does. The matrix

$$H_2 = \begin{bmatrix} \alpha_0 & \cdots & \alpha_{n-1} \\ \alpha_0^3 & \cdots & \alpha_{n-1}^3 \end{bmatrix}$$

is the parity-check matrix of a binary code of length $n = 2^m - 1$ which corrects up to two errors. A vector $\mathbf{c} = (\, c_0 \quad \cdots \quad c_{n-1}\,)$ in $V_n(F_2)$ is a code word in the code defined by H_2 if and only if $\sum_{i=0}^{n} c_i \alpha_i = \sum_{i=0}^{n} c_i \alpha_i^3 = 0$. Since the $2m$ rows of the matrix H_2 over F_2 may not be all linearly independent, the dimension of the code is $d(C) \geq n - 2m = 2^m - 1 - 2m$.

§

Definition 13. The *Vandermonde matrix* is defined as

$$A = \begin{bmatrix} 1 & \cdots & 1 \\ a_1 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_1^{r-1} & \cdots & a_r^{r-1} \end{bmatrix}$$

§

Theorem 9. Let $a_1, \ldots, a_r$ be distinct non-zero elements of a field. Then the the Vandermonde matrix is such that

$$\begin{vmatrix} [1 & \cdots & 1 \\ a_1 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_1^{r-1} & \cdots & a_r^{r-1} \end{vmatrix} \neq 0$$

Proof. Subtracting row$(i+1) - a_1$ row i, $i = 1, \cdots, r-1$, yields,

$$\begin{aligned} \det A &= \begin{vmatrix} 1 & 1 & \cdots & 1 \\ 0 & a_2 - a_1 & \cdots & a_r - a_1 \\ 0 & a_2(a_2 - a_1) & \cdots & a_r(a_r - a_1) \\ \vdots & & \ddots & \vdots \\ 0 & a_2^{r-2}(a_2 - a_1) & \cdots & a_r^{r-2}(a_r - a_1) \end{vmatrix} \\ &= (a_2 - a_1) \cdots (a_r - a_1) \begin{vmatrix} 1 & \cdots & 1 \\ a_2 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_2^{r-2} & \cdots & a_r^{r-2} \end{vmatrix} \\ &= (a_2 - a_1) \cdots (a_r - a_1) \cdot (a_3 - a_2) \cdots (a_r - a_2) \begin{vmatrix} 1 & \cdots & 1 \\ a_3 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_3^{r-3} & \cdots & a_r^{r-3} \end{vmatrix} \\ &\vdots \\ &= \prod_{i>j} (a_i - a_j) \end{aligned}$$

Then, since a_i are distinct and non-zero, therefore det A is non-zero. ¶

Theorem 10. Any square matrix having a non-zero determinant has all its columns linearly independent.

Proof. Let A be an $r \times r$ matrix, and that $|\,A\,| \neq 0$. Then suppose the columns of A are linearly dependent. Then one may write some column of A as a linear combination of the others, for example

$$\mathbf{c}_j = \sum_{\substack{i=1 \\ i \neq j}}^{r} a_i \mathbf{c}_i$$

Then if column $\mathbf{c}_j$ is replaced by $\mathbf{c}_j - \sum_{\substack{i=1 \\ i \neq j}}^{r} a_i \mathbf{c}_i$ gives a matrix B with $|\,B\,| = |\,A\,|$. But B also has a column whose all elements are zeros, which means that $|\,A\,| = |\,B\,| = 0$, a contradiction and thus the proof. ¶

Theorem 11. Let $(\alpha_0, \ldots, \alpha_{n-1})$ be an ordering of non-zero elements of $\mathbf{F}_{2^m}$, and let t be a positive integer such that $t \leq 2^{m-1} - 1$. Then the matrix

$$H = \begin{bmatrix} \alpha_0 & \cdots & \alpha_{n-1} \\ \alpha_0^3 & \cdots & \alpha_{n-1}^3 \\ \vdots & \ddots & \vdots \\ \alpha_0^{2t-1} & \cdots & \alpha_{n-1}^{2t-1} \end{bmatrix}$$

is the parity-check matrix of a binary (n, k)-code capable of correcting all error patterns of weight $w \leq t$, with dimension $k \geq n - mt$.

Proof. A vector $\mathbf{c} = (c_0, \ldots, c_{n-1})$ in $V_n(F_2)$ is a code word if and only if $H\mathbf{c}^T = \mathbf{0}$. Thus,

$$\sum_{i=0}^{n-1} c_i \alpha_i^j = 0$$

for $j = 1, 3, \ldots, 2t-1$. We simplify this by using the fact that $(x+y)^2 = x^2 + y^2$ in characteristic 2, and $x^2 = x$ in F_2. Hence,

$$\left(\sum_{i=0}^{n-1} c_i \alpha_i^j \right)^2 = \sum_{i=0}^{n-1} c_i^2 \alpha_i^{2j} = \sum_{i=0}^{n-1} c_i \alpha_i^{2j}$$

for $j = 1, 3, \ldots, 2t-1$, which gives us

$$\sum_{i=0}^{n-1} c_i \alpha_i^j$$

for $j = 1, 2, \ldots, 2t$. Therefore we could also use the parity-check matrix

$$H' = \begin{bmatrix} \alpha_0 & \cdots & \alpha_{n-1} \\ \alpha_0^2 & \cdots & \alpha_{n-1}^3 \\ \vdots & \ddots & \vdots \\ \alpha_0^{2t} & \cdots & \alpha_{n-1}^{2t} \end{bmatrix}$$

According to Theorem 8 H' is a parity-check matrix which corrects t errors if and only if every subset of $2t$ or fewer columns of H' is linearly independent. Next, since a subset of $r \leq 2t$ columns of H' has the form

$$A = \begin{bmatrix} a_1 & \cdots & a_r \\ a_1^2 & \cdots & a_r^2 \\ \vdots & \ddots & \vdots \\ a_1^{2t} & \cdots & a_r^{2t} \end{bmatrix}$$

where $a_1, \ldots, a_r$ are distinct non-zero elements of F_{2m}, we may consider the matrix

$$A' = \begin{bmatrix} a_1 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_1^r & \cdots & a_r^r \end{bmatrix}$$

which is nonsingular since its determinant by the Vandermonde determinant theorem, Theorem 9, is

$$\det A' = a_1 \cdots a_r \begin{vmatrix} 1 & \cdots & 1 \\ a_1 & \cdots & a_r \\ \vdots & \ddots & \vdots \\ a_1^{r-1} & \cdots & a_r^{r-1} \end{vmatrix} = a_1 \cdots a_r \prod_{i<j} (a_j - a_i) \neq 0$$

Then the columns of A', and hence those of A, cannot be linearly dependent, and therefore the code corrects all error patterns of weight up to t. Now H, as a matrix with entries from F_2 rather than F_{2m}, has dimensions $mt \times n$, hence the dual code has dimension $k \leq mt$, and the code has dimension $k \geq n - mt$. ¶

Theorem 12. Let C be a linear (n,k)-code over $GF(q)$ with parity-check matrix H. Then the minimum distance of C is d if and only if any $d-1$ columns of H are linearly independent but some d columns are linearly dependent.

Proof. The minimum distance of a code $d(C)$ is equal to the minimum of the weights of the non-zero code words. Let $\mathbf{x} = x_1 \cdots x_n$ be a vector in $V(n,q)$. Then $\mathbf{x}$ is in C if and only if $\mathbf{x}H^T = \mathbf{0}$ if and only if $x_1\mathbf{h}_1 + \cdots + x_n\mathbf{h}_n = \mathbf{0}$, where $\mathbf{h}_1, \ldots, \mathbf{h}_n$ are the columns of H. Therefore there is a set of d linearly dependent columns of H corresponding to each code word $\mathbf{x}$ of weight d. On the other hand, if there existed a set of $d-1$ linearly dependent columns of H, then there would exist some scalars $x_{i_1}, \ldots, x_{i_{d-1}}$, not all zero, such that $\sum_{j=1}^{d-1} x_{i_j} = \mathbf{0}$. But if this were the case, then $\mathbf{x}H^T = \mathbf{0}$ and so would be a code word of weight $0 < d < d(C)$. ¶

Theorem 13. The maximum dictionary size m such that there exists a q-ary (n,m,d)-code is $A_q(n,d) \leq q^{n-d+1}$.

Proof. Let C be a q-ary (n,m,d)-code. If we remove the last $d-1$ coordinates from each code word, then the m vectors of length $n-d+1$ so obtained must be distinct, otherwise $d(C)$ must be less than d, which would contradict the statement above. Therefore $m \leq q^{n-d+1}$. ¶

Theorem 14. Let C be the code over $GF(q)$, where q is a prime number, and C is defined to have the parity-check matrix

$$H = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & 2 & 3 & \cdots & n \\ 1 & 2^2 & 3^2 & \cdots & n^2 \\ \vdots & & & \ddots & \vdots \\ 1 & 2^{d-2} & 3^{d-2} & \cdots & n^{d-2} \end{bmatrix}$$

where $d \leq n \leq q-1$. If q is a prime-power, then $A_q(n,d) = q^{n-d+1}$.

Proof. We have,

$$C = \left\{ x_1 \cdots x_n \in V(n,q) \text{ s.t. } \sum_{i=1}^{n} i^j x_i = 0 \text{ for } j = 0, 1, \ldots, d-2 \right\}$$

Any $d-1$ columns form a Vandermonde matrix, and therefore by Theorem's 9 and 10 are linearly independent. By Theorem 12 C has a minimum distance d and therefore is a q-ary (n, q^{n-d+1}, d)-code. The proof follows since C meets the Singleton bound of Theorem 13. ¶

Problem 4. Find the decoding procedure for the BCH codes.

Solution. Assume that $d = 2t+1$ and H has $2t$ rows. Suppose the code word $\mathbf{c} = c_1 \cdots c_n$ is transmitted and the vector $\mathbf{r} = r_1 \cdots r_n$ is received. Assuming that at most t errors have occurred, let $x_1, \ldots, x_t$ be their positions and $m_1, \ldots, m_t$ their respective magnitudes. Then the syndrome is

$$(s_1, \ldots, s_{2t}) = \mathbf{r}H^T$$

and we have

$$s_j = \sum_{i=1}^{n} r_i i^{j-1} = \sum_{i=1}^{t} m_i x_i^{j-1} \tag{3}$$

for $j = 1, \ldots, 2t$. Then from

$$\phi(\theta) = \frac{m_1}{1 - x_1\theta} + \frac{m_2}{1 - x_2\theta} + \cdots + \frac{m_t}{1 - x_t\theta} \tag{4}$$

and

$$\frac{m_i}{1 - x_i\theta} = m_i \left(1 + x_i\theta + x_i^2\theta^2 \cdots\right)$$

together with Equation 3, we have

$$\phi(\theta) = s_1 + s_2\theta + \cdots + s_{2t}\theta^{2t-1} + \cdots$$

Also, from Equation 4 we have

$$\phi(\theta) = \frac{a_1 + a_2\theta + a_3\theta^2 + \cdots + a_t\theta^{t-1}}{1 + b_1\theta + b_2\theta^2 + \cdots + b^t\theta^t} \tag{5}$$

Hence,

$$\left(s_1 + s_2\theta + s_3\theta^2 + \cdots\right)\left(1 + b_1\theta + b_2\theta^2 + \cdots + b_t\theta^t\right) = a_1 + a_2\theta + \cdots + a_t\theta^{t-1}$$

Which gives us

$$a_1 = s_1 \quad \text{and} \quad a_i = \sum_{j=0}^{i-1} s_{i-j} b_i, \quad i = 2, \ldots, t \tag{6}$$

and

$$0 = \sum_{j=0}^{t} s_{i-j} b_j, \quad i = t+1, \ldots, 2t \tag{7}$$

With a_i and b_i known we may turn Equation 5 into partial fractions

$$\phi(\theta) = \frac{p_1}{1 - q_1\theta} + \cdots + \frac{p_t}{1 - q_t\theta}$$

and therefore $m_i = p_i$ and $x_i = q_i$, for $i = 1, \ldots, t$, and the system in Equation 3 is solved. Algorithm 1 then gives the procedure for error correction.

#

Note 7. The polynomial

$$\sigma(\theta) = 1 + b_1\theta + b_2\theta^2 + \cdots + b_t\theta^t = (1 - x_1\theta)\cdots(1 - x_t\theta) \tag{8}$$

can be used to locate the location of the errors. The polynomial

$$\omega(\theta) = a_1 + a_2\theta + \cdots + a_t\theta^{t-1}$$

can be used to find the magnitude of the errors.

§

Algorithm 1 *Procedure for correcting up to t errors in BCH codes.*

input: **r**
find $s_1, \ldots, s_2t$
$e \leftarrow$maximum number of equations in Equation 7
for $i = e + 1$ to t **do**
  $b_i \leftarrow 0$
endfor
$(b_1, \ldots, b_e) \leftarrow$**solve** the first e equations of Equation 7
$(z_1, \ldots, z_e) \leftarrow$**find** the e zeros of Equation 8
$(a_1, \ldots, a_e) \leftarrow$**solve** Equation 6
for $i = 1$ to e **do**
  $m_i \leftarrow \frac{a_1 + a_2x_i + \cdots + a_ex_i^{e-1}}{\prod_{\substack{j=1 \\ j\neq i}}^{e}(1 + x_jx_i)}$
endfor

Bibliography

Raymond Hill. *A first course in coding theory.* Clarendon, 1986
San Ling and Chaoping Xing. *Coding theory, a first course.* Cambridge University Press, 2004
Robert J McEliece *The theory of information and coding.* Addison-Wesley, 1977
George F Simmons. *Topology and modern analysis.* McGraw-Hill, 1963
L R Vermani. *Elements of algebraic codng theory.* Chapman & Hall, 1996